

LAX PAIR FORMULATION OF THE W-GRAVITY THEORIES IN TWO DIMENSIONS

Jean-Marc Lina and Prasanta K. Panigrahi

*Laboratoire de Physique Nucléaire
Université de Montréal
C.P. 6128, succ "A"
Montréal, Québec, Canada
H3C 3J7*

Abstract

The Lax pair formulation of the two dimensional induced gravity in the light-cone gauge is extended to the more general w_N theories. After presenting the w_2 and w_3 gravities, we give a general prescription for an arbitrary w_N case. This is further illustrated with the w_4 gravity to point out some peculiarities. The constraints and the possible presence of the cosmological constants are systematically exhibited in the zero-curvature condition, which also yields the relevant Ward identities. The restrictions on the gauge parameters in presence of the constraints are also pointed out and are contrasted with those of the ordinary 2d-gravity.

Submitted to International Journal of Modern Physics A

I. Introduction

Induced gravity in two dimensions, arising from the interaction of conformal matter with the gravitational field, is now being extensively studied, due mainly to its relevance to string theory^[1]. Significant progress has been made in the weak-coupling regime ($c \leq 1$ and $c > 25$, c being the central charge of the matter sector), both in the direct continuum^[2] and the lattice model approaches^[3]. Although the non-local effective action, given by

$$S = \alpha \int \sqrt{-g} (R \square^{-1} R - \lambda) d^2 x, \quad (1)$$

becomes local both in the conformal and the light-cone gauges, the problem of quantization is much more tractable in the latter case, as was first pointed out by Polyakov^[1]. In particular, a hidden $sl(2, R)$ Kac-Moody symmetry becomes apparent in this approach, allowing the extraction of non-perturbative informations about the theory, e.g the anomalous dimensions of various fields in the presence of quantum gravitational fields^[4]. The light-cone gauge is characterised by the line element

$$ds^2 = dx^+ dx^- + h(x^+, x^-) dx^{+2}, \quad (2)$$

where the metric $h(x^+, x^-)$ transforms under the residual coordinate transformations as

$$\delta h = \epsilon h' + \dot{\epsilon} - h \epsilon'. \quad (3)$$

The dot and the prime denote the derivatives with respect to x^+ and x^- variables. The simplicity of the light-cone approach can be easily seen from the equation of constant curvature, which yields a constraint

$$R = \partial_-^2 h = -2\lambda, \quad (4)$$

in contrast to the dynamical Liouville equation in the conformal gauge. The constraint equation yields the equation of motion for the induced gravity $\partial_-^3 h = 0$, allowing the expansion of $h(x^+, x^-)$ as

$$h(x^+, x^-) = J^-(x^+) - 2J^0(x^+)x^- + J^+(x^+)x^{-2}; \quad (5)$$

here $J^-(x^+)$, $J^0(x^+)$ and $J^+(x^+)$ are the currents of the $sl(2, R)$ Kac-Moody algebra.

A particularly clear explanation of the geometrical origin of the current algebra symmetries in the induced gravity was provided by Polyakov^[5], who demonstrated how to get

diffeomorphisms from the restricted gauge transformations. Considering a $sl(2, R)$ Lie algebra valued field $A_\mu dx^\mu$, a partial gauge fixing in the A_- sector led to the transformation of the A_-^+ field as the stress-energy tensor of a conformal field theory. Suitable gauge fixing in the A_+ sector then led to the transformation of the A_+^- component as the light-cone metric; surprisingly, the gravitational Ward identity emerged as a consistency condition of the gauge fixing or, equivalently, from the zero-curvature condition. Generalizations to the $sl(N, R)$ case are now being actively pursued in the literature^[6].

In the works of Das et al.^[7], the zero-curvature condition, written as a compatibility equation of a matrix Lax pair, was interpreted as a gauge anomaly equation. The anomaly equation can, in principle, be integrated to yield a suitably gauged WZWN action^[8]. Following this approach, the KdV and the Boussinesq hierarchies were related with the ordinary and the w_3 gravitational Ward identities respectively. Subsequently, the work of the present authors incorporated the curvature constraints in this Lax pair formulation of the 2-d gravity^[9].

In the present work, our goal is to analyse the more general w_N gravities. In particular, we will provide a general prescription to determine the A_- and the A_+ sectors of the $sl(N, R)$ Lie algebra valued gauge connection such that the compatibility of the Lax pair, $\partial_- + A_-$ and $\partial_+ + A_+$, will yield the constraints and the dynamical equations. We will briefly explain the origin of the constraints in the w_N -gravity, and their relevance for the quantum regime of the theory. The elements of the A_- field, w_2 and $w_{k, k>2}$, will transform respectively as the stress energy tensor and the spin- k quasi-primary fields under the residual gauge transformations. A conjectured formula, valid to all orders, will be provided for the infinitesimal variations of these fields under diffeomorphism. The A_+ sector contains variables which transform as the currents of a conformal field theory. These fields will be interpreted as the metrics of the w_N gravity^[6,10].

In Sec. II, we give a brief introduction to the w_N algebras and their connection with the integrable non-linear equations. Our approach is outlined in the well studied $sl(2, R)$ and $sl(3, R)$ cases in Sec. III. Sec. IV deals with the generalisation to an arbitrary w_N gravity. As an illustration, the w_4 gravity is also investigated in this section, pointing out certain peculiarities in this case and showing explicitly how some non-linear evolution equations can arise as special cases of the w_4 Ward identities. We conclude in Sec.V with some final remarks and future directions of work. There are three appendices; the first one gives the variations of the fields in $sl(4, R)$, the second lists the variations of the w_i 's in $sl(10, R)$ as a specific example of the conjectured general formula given in Sec. IV. The last appendix presents the infinitesimal variation of the metric field h_2 in some higher spin w -gravity theories.

II. w_N algebras and covariant Lax operators

A generalization of the Virasoro algebra to the so-called w_3 algebra¹ was achieved by Zamolodchikov^[11]. This algebra contains a chiral spin-3 conserved current $w_3(z)$ in addition to the stress-energy tensor $w_2(z)$. With additional fields, consistent generalizations to higher spin algebras have been discovered in the literature^[12] and these w_N algebras have appeared in many physical contexts. To quote a few examples, they have manifested in the gauged WZWN models^[13], in the cosets of affine Lie algebras^[14], Toda field theories^[15], $2 + 1$ dimensional Chern-Simons theories^[16] and also in the context of various non-linear integrable equations^[17] e.g. KdV, Boussinesq, etc. It is worth emphasizing that these algebras are not Lie algebras because of quadratic defining relations as will become clear in the course of the text. The w_N algebras have series of unitary representations characterised by

$$c = (n-1) \left(1 - \frac{n(n-1)}{p(p+1)} \right), \quad p = n+1, n+2, \dots \quad (6)$$

and hence have been of interest in the construction of the “periodic table” of the conformal-invariant solutions of the two-dimensional euclidean quantum field theories. Here, we will be mainly concerned with the classical w_N algebras. In particular, their connection with the non-linear integrable equations is pertinent for our work.

The much studied KdV equation

$$w_{2t} = w_{2xxx} + 6w_2w_{2x}, \quad (\text{where } w_{2x} \equiv \frac{\partial w_2}{\partial x}), \quad (7)$$

and the related hierarchy of equations can be written as Hamiltonian equations

$$w_{2t} = \{w_2, \mathcal{H}_n\}_{(l)} \quad (8)$$

in two distinct ways^[18]. The two Hamiltonians

$$\mathcal{H}_1 = \frac{1}{2} \int dx (2w_2^3 - w_{2x}^2), \quad \text{and} \quad \mathcal{H}_2 = \int dx w_2^2 \quad (9)$$

give rise to the above equation if the respective Poisson brackets ($l = n$) are defined as

$$\{w_2(x), w_2(y)\}_{(1)} = \partial \delta(x-y), \quad \text{and} \quad \{w_2(x), w_2(y)\}_{(2)} = \left(\frac{1}{2} \partial^3 + w_2 \partial + \partial w_2 \right) \delta(x-y) \quad (9)$$

where the operator $\partial = \frac{\partial}{\partial x_-}$ acts on all the objects to its right. The two Poisson brackets can be recognized as the abelian current algebra and the Virasoro algebra respectively.

¹ In the present day terminology, this algebra is known as the $w_3^{(1)}$ algebra.

In Ref.[19], another interesting derivation of the KdV equation, connecting it to the covariance property of the Hill operator under reparameterisation was obtained. It has been subsequently extended to other integrable systems^[20]. Briefly, the Hill operator

$$\mathcal{L}_{(2)} = \partial^2 + w_2(x^-) \quad (11)$$

transforms covariantly under reparameterisation in the sense that

$$\begin{aligned} x^- &\rightarrow f(x^-), \\ \mathcal{L}_{(2)} &\rightarrow (f')^{\frac{-3}{2}} (\partial^2 + \overline{w}_2(x^-)) (f')^{\frac{-1}{2}} \end{aligned} \quad (12)$$

where

$$\overline{w}_2(x^-) = (f')^2 w_2(f) + \frac{1}{2} S_f(x^-),$$

S_f being the Schwarzian derivative, $S_f = \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'} \right)^2$. A time (represented by x^+) dependent reparameterisation then yields $w_2(x^-) \rightarrow \overline{w}_2(x^+, x^-)$ and

$$\dot{\overline{w}}_2 = \frac{1}{2} h''' + 2\overline{w}_2 h' + \overline{w}_2' h, \quad (13)$$

where h is defined as $\frac{\dot{f}}{f'}$. The KdV equation in the standard form is obtained by making the special choice $h = 2\overline{w}_2$. It is worth pointing out that the above non-linear evolution equation is nothing but the gravitational Ward identity for the central charge $c = \frac{1}{2}$ and $h = \frac{\dot{f}}{f'}$ is the well known Beltrami equation defining the light-cone metric.

At this point, looking at the infinitesimal variation of $w_2(x^-)$ under $x^- \rightarrow x^- + \epsilon^-$, the second Poisson bracket mentioned before can be extracted, assuming $w_2(x^-)$ as the generator of diffeomorphism.

This procedure, in principle, can be generalized to the higher covariant operators. Given an n th order differential operator of the form $M_n = \partial^n + \sum_{i=0}^{n-2} u_i \partial^i$, a covariant operator $\mathcal{L}_{(n)}$ can be constructed^[21] such that it acts on densities of weight $\frac{1-n}{2}$ and the functions appearing in $\mathcal{L}_{(n)}$ are the generators of the w_N algebra. Some examples of $\mathcal{L}_{(n)}$ are given below.

$$\begin{aligned} \mathcal{L}_{(0)} &= 1, \\ \mathcal{L}_{(1)} &= \partial, \\ \mathcal{L}_{(2)} &= \partial^2 + w_2, \\ \mathcal{L}_{(3)} &= \partial^3 + 4w_2 \partial + 2w_{2x} + w_3, \\ \mathcal{L}_{(4)} &= \partial^4 + 10w_2 \partial^2 + 10w_{2x} \partial + 9w_2^2 + 3w_{2xx} + \partial w_3 + w_3 \partial + w_4. \end{aligned} \quad (14)$$

It should be mentioned that the covariant Lax operators $\mathcal{L}_{(2)}$ and $\mathcal{L}_{(3)}$ have appeared in the w_2 and w_3 gravities, in the works of Zamolodchikov^[22] and Matsuo^[23] respectively.

For the sake of completeness, it could also be pointed out that the above mentioned Hamiltonian formulations can be related to the symplectic structures associated with the space of pseudo-differential operators^[24]. In particular, the second Poisson bracket structure relevant for the w_N algebras can be extracted following Adler^[25].

Given a Lax operator $\mathcal{L}_{(n)}$, one defines

$$\frac{\partial \mathcal{L}_{(n)}}{\partial t} = (\mathcal{L}_{(n)}F)_+ \mathcal{L}_{(n)} - \mathcal{L}_{(n)}(\mathcal{L}_{(n)}F)_+ \quad (15)$$

where

$$F = \partial^{-1}f_0 + \partial^{-2}f_1 + \dots + \partial^{-n}f_{n-1},$$

and the $(\mathcal{L}_{(n)}F)_+$ is the differential part of $\mathcal{L}_{(n)}F$. Comparing this equation with

$$\frac{\partial u_i}{\partial t} = \sum_{j=0}^{n-2} D_{ij}^{(2)} f_j \quad (16)$$

one finds the Poisson brackets as $\{u_i(x), u_j(y)\} = D_{ij}^{(2)} \delta(x - y)$. The Lax operators, if written in a covariant form, generate the w_N algebras.

Although, we will pursue a method involving the matrix valued Lax pairs in this paper, it turns out that the specific mapping from M_n to $\mathcal{L}_{(n)}$ is of relevance in this formalism. The relationship of the u_i 's and the w_i 's appearing in M_n and $\mathcal{L}_{(n)}$ respectively is precisely those of the fields W_i 's and the primary fields w_i 's in this work. The scalar differential operators will be connected with the matrix Lax operator $\partial_- + A_-$ to explain this interesting relationship. It is also worth observing that the inverse mapping appears to determine the A_+ sector and the gauge parameters that generate the gauge preserving transformations.

III. w_2 and w_3 gravities

Let us first elaborate on the (matrix) Lax pair formulation of the much studied ordinary gravity. This is done in a manner which can be easily generalized to an arbitrary w_N algebra. In case of the w_2 gravity, one considers the deformed $sl(2, R)$ algebra^[26]

$$[\mathbf{t}_+, \mathbf{t}_-] = \lambda_+ \lambda_- \mathbf{t}_0, \quad [\mathbf{t}_0, \mathbf{t}_\pm] = \pm 2\mathbf{t}_\pm. \quad (17)$$

where $\mathbf{t}_a$'s are represented by the matrices

$$\mathbf{t}_- = \begin{pmatrix} 0 & 0 \\ \lambda_- & 0 \end{pmatrix}, \quad \mathbf{t}_+ = \begin{pmatrix} 0 & \lambda_+ \\ 0 & 0 \end{pmatrix}, \quad \mathbf{t}_0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}; \quad (18)$$

$\lambda_{\pm}$ being the so-called spectral parameters. In the limit of $\lambda_{\pm}$ going to zero, the algebra reduces to the Poincaré algebra in two dimensions. The Killing metric $g_{ab} = \frac{1}{2}f_{ac}{}^d f_{bd}{}^c$ is degenerate when either λ_+ or λ_- is vanishing. Considering the reduced Killing metric on $sl(2, R)/U(1)$ given by $2\lambda_+\lambda_- \eta_{ab}$, we define the gauge field $A = A^a t_a$ and the line element as the coset invariant

$$ds^2 = \eta_{ab} A^a A^b. \quad (19)$$

The gauge fields are the zweibeins and the spin connection.

The following chiral gauge choice

$$A_- = \begin{pmatrix} 0 & 0 \\ \lambda_- & 0 \end{pmatrix}, \quad A_+ = \begin{pmatrix} -\frac{1}{2}h' & \lambda_+ \\ \lambda_- h & \frac{1}{2}h' \end{pmatrix}, \quad (20)$$

yields the constraint equation

$$R = \partial^2 h = -2\lambda_+\lambda_-$$

as the integrability condition, $F = dA + A \wedge A = 0$. It can be easily checked that the field $h(x^+, x^-)$ transforms as the light-cone metric. Furthermore, it is worth pointing out that the same equations can be obtained as a consistency condition under the gauge transformations that maintain this gauge choice. Following the remarkable analysis of Polyakov^[5], we can show that the gauge choice

$$A_- = \begin{pmatrix} 0 & -t \\ \lambda_- & 0 \end{pmatrix}, \quad A_+ = \begin{pmatrix} -\frac{1}{2}h' & \lambda_+ - (\frac{1}{2\lambda_-} + \kappa)h'' - th \\ \lambda_- h & \frac{1}{2}h' \end{pmatrix} \quad (21)$$

yields $\delta t = 2t\epsilon' + \epsilon t' + \frac{1}{2\lambda_-}\epsilon'''$ under the residual gauge transformations that maintain A_- . This variation is easily recognized as the variation of the stress-energy tensor in a conformal field theory, with a central charge c equal to $\frac{1}{2\lambda_-}$. Furthermore, all the parameters of the gauge variation are not independent:

$$\begin{cases} \epsilon^- = +\frac{1}{\lambda_-}\epsilon, \\ \epsilon^0 = -\frac{1}{2}\partial\epsilon, \\ \epsilon^+ = -(\frac{1}{2\lambda_-}\partial^2 + t)\epsilon; \end{cases} \quad (22)$$

and it can be easily checked that $h(x^+, x^-)$ does transform as the light-cone metric field:

$$\delta h = -h\epsilon' + \epsilon h' + \dot{\epsilon}. \quad (23)$$

The consistency requirement for maintaining the gauge choice or the zero curvature condition $F = 0$ provide the constraint equation and the gravitational Ward identity given respectively by

$$\partial^2 h = \frac{\lambda_+}{\kappa}, \quad (24a)$$

and

$$\partial_+ t = \left((c + \kappa) \partial_-^3 + \partial t + t \partial \right) h. \quad (24b)$$

The work of Das et al.^[7] corresponds to the case $\kappa = \lambda_+ = 0$, for which only the Ward identity exists. It is worth emphasising that for taking into account the constraint with the possible presence of a cosmological constant, the parameter κ should be nonvanishing. It should be pointed out that it is precisely the case in the quantum Ward identity of the 2d-gravity. In this case however, the nonvanishing κ is exactly calculable. In our gauge choice with $\kappa \neq 0$, the gauge preserving symmetries are restricted to those described by ϵ such that $\partial \epsilon = 0$ if $\lambda_+ \neq 0$ or $\partial^2 \epsilon = 0$ if $\lambda_+ = 0$. The parameter of the residual coordinate invariance is restricted as can be observed in the work of KPZ^[4].

We pursue the analysis with the example of $N = 3$ for the purpose of generalization to an arbitrary w_N case. We choose the following matrix representation for the $sl(3, R)$ algebra elements (the same as in Ref.[7]):

$$\begin{aligned} t_0 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad t_+ = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad t_- = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ t_{00} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad t_{++} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad t_{--} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \\ t_{0+} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad t_{0-} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \end{aligned} \quad (25)$$

and define the gauge field A_- :

$$A_- = \begin{pmatrix} 0 & -W_2 & -W_3 \\ c_1 & 0 & 0 \\ 0 & c_2 & 0 \end{pmatrix} \quad (26)$$

where the c_i 's are constants and the W_i 's are dynamical fields with dimension i (the dimension of $\frac{\partial}{\partial x_-}$ is $+1$). As we have noticed in Ref.[9], one can set $c_2 = 1$ without loss of generality².

As they stand, the dynamical fields in A_- do not satisfy the Virasoro symmetry properties mentioned before. The gauge transformations that preserve Eq.(26) are found to be

² This constant scales the primary field w_3 defined in the next page; this observation generalizes to the higher spin algebras.

described by the functions ϵ^- and ϵ^{--} or, more conveniently, by the arbitrary functions ρ_2 and ρ_3 :

$$\begin{cases} \epsilon^- = c_1 \rho_2 - \frac{c_1}{2} \rho'_3, \\ \epsilon^{--} = c_1 \rho_3, \end{cases} \quad (27)$$

and, defining the w_i 's by the following change of variables

$$W_2 = w_2, \quad W_3 = w_3 - \frac{1}{2} w'_2, \quad (28)$$

$$\begin{cases} \epsilon^0 = -\rho'_2 + c_1 \left(\frac{1}{6c_1} \partial^2 - \frac{1}{3} w_2 \right) \rho_3, \\ \epsilon^+ = -\left(\frac{1}{c_1} \partial^2 + w_2 \right) \rho_2 + \left(\frac{1}{6c_1} \partial^3 + \frac{1}{6} \partial w_2 - w_3 \right) \rho_3, \\ \epsilon^{00} = -\rho'_2 - c_1 \left(\frac{1}{6c_1} \partial^2 + \frac{2}{3} w_2 \right) \rho_3, \\ \epsilon^{0-} = \rho_2 + \frac{1}{2} \rho'_3, \\ \epsilon^{0+} = -\rho''_2 - c_1 \left(\frac{1}{6c_1} \partial^3 - \frac{1}{6} \partial w_2 - \frac{1}{2} w_2 \partial - w_3 \right) \rho_3. \end{cases} \quad (29)$$

Indeed, the gauge preserving transformations lead to the Virasoro and the spin-3 symmetries in terms of the parameters ρ_2 and ρ_3 as defined in Eq.(27). These symmetries are generated respectively by the energy momentum tensor w_2 and the primary field w_3 defined in Eq.(28) since we find that $\delta_\epsilon(A_-)$ leads to

$$\delta w_i = \delta(\rho_2) w_i + \delta(\rho_3) w_i$$

with

$$\begin{aligned} \delta(\rho_2) w_2 &= 2w_2 \rho'_2 + \rho_2 w'_2 + \frac{2}{c_1} \rho_2''', \\ \delta(\rho_2) w_3 &= 3w_3 \rho'_2 + \rho_2 w'_3, \\ \delta(\rho_3) w_2 &= 3w_3 \rho'_3 + 2\rho_3 w'_3, \\ \delta(\rho_3) w_3 &= -\frac{1}{12} (2w_2''' \rho_3 + 9w_2'' \rho'_3 + 15w_2' \rho_3'' + 10w_2 \rho_3''') \\ &\quad - \frac{2c_1}{3} (w_2^2 \rho'_3 + w_2 w'_2 \rho_3) - \frac{1}{6c_1} \rho_3'''''. \end{aligned} \quad (30)$$

We can consistently construct a Poisson bracket structure between the w_i 's, starting with the Virasoro algebra satisfied by the generator of diffeomorphisms w_2 :

$$\{w_2(x), w_2(y)\} = (c\partial^3 + w_2\partial + \partial w_2)\delta(x-y) \quad (31)$$

with $c = \frac{2}{c_1}$. The primary field w_3 can be interpreted as the generator of the spin-3 transformations:

$$\begin{aligned}\{w_3(x), w_3(y)\} &= -\frac{1}{12}(c\partial^5 + 2w_2\partial^3 + 3\partial w_2\partial^2 + 3\partial^2 w_2\partial + 2\partial^3 w_2 + \frac{16}{c}w_2\partial w_2)\delta(x-y), \\ \{w_2(x), w_3(y)\} &= (w_3\partial + 2\partial w_3)\delta(x-y) \\ \{w_3(x), w_2(y)\} &= (2w_3\partial + \partial w_3)\delta(x-y).\end{aligned}\tag{32}$$

As can be seen from the Poisson bracket $\{w_3(x), w_3(y)\}$, there are non-linear terms which go to zero in the so-called “classical limit” $c \rightarrow \infty$. The A_- sector being defined,

$$A_- = \begin{pmatrix} 0 & -w_2 & -w_3 + \frac{1}{2}w_2' \\ c_1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \tag{33}$$

we now study the A_+ gauge field. The components A_+^a are defined in terms of the “metric fields” h_i , $i = 2, 3$ and some general n th order differential operator, *linear* in the W_i ’s and denoted by $\mathcal{D}_a^{(n)}$:

$$A_+^a = \mathcal{D}_a^{(1+l(a))}h_2 + \mathcal{D}_a^{(2+l(a))}h_3. \tag{34}$$

Here, $l(a)$ represents the number of nonvanishing gauge indices in “ a ” e.g. $l(+) = l(0+) = 1$, $l(-) = l(0-) = -1$, $l(0) = l(00) = 0$, and $l(++) = -l(--) = 2$. Finally, we introduce the possible “cosmological constants” λ_i through the shift

$$A_+ \rightarrow A_+ + \lambda_2 \mathbf{t}_+ + \lambda_3 \mathbf{t}_{0+}, \tag{35}$$

a choice which will be justified shortly. Let us therefore write the gauge field in the form

$$A_+ = \begin{pmatrix} \dots & \lambda_0 + \dots & \dots \\ H_2 & \dots & \lambda_1 + \dots \\ H_3 & \dots & \dots \end{pmatrix}, \tag{36}$$

where we have emphasized the components $H_i \sim h_i + \dots$, i.e. the components A_+^- and A_+^{--} . Notice that the gauge fields are, by construction, *linear* in the fields W_i ’s and the prescription used to describe them is easily generalizable to higher rank $sl(N, R)$ algebras. The significance of the constants is already known^[9] for the $sl(2, R)$ and the $sl(3, R)$ cases: the λ_i ’s enter in the constraint equations involving the h_i ’s, and, for the w_2 -gravity, λ_2 can be interpreted as the cosmological constant.

The components of A_+ are determined in such a way that the curvature is of the following form

$$\begin{pmatrix} \Phi_2 & E_1 & E_2 \\ 0 & \Phi_3 - \Phi_2 & 0 \\ 0 & 0 & -\Phi_3 \end{pmatrix}, \tag{37}$$

with

$$\begin{cases} \Phi_2 = \kappa_2 \partial^2 h_2 - \lambda_2, \\ \Phi_3 = \kappa_3 \partial^3 h_3 - \lambda_3 \end{cases}, \quad (38)$$

and $E_1 = -\partial_+ w_2 + \dots$, $E_2 = -\partial_+ w_3 + \dots$. The constants κ_i are free parameters. Together with the “cosmological constants” λ_i , they represent the presence or the absence of the constraints. Let us notice that the Eqs.(38) are legitimate only if the metric fields h_i have the proper dimensions such that $\partial^i h_i$ is dimensionless. In fact, the zero-curvature condition *alone* does not define a unique A_+ since we obtain

$$\begin{cases} H_2 = c_1 h_2 + a h'_3, \\ H_3 = c_1 h_3, \end{cases} \quad (39)$$

where a is free. However, choosing this parameter equal to zero spoils the symmetry property of h_2 under the diffeomorphisms. These symmetries are easily calculated from Eqs.(26,28) and $\delta A_+ = \partial_+ \epsilon + [A_+, \epsilon]$. Keeping the interpretation of h_2 as a metric field, i.e. Eq.(3), gives $a = -\frac{c_1}{2}$. Then, the remaining variations are found to be

$$\begin{aligned} \delta(\rho_2) h_3 &= -2h_3 \rho'_2 + \rho_2 h'_3, \\ \delta(\rho_3) h_2 &= \frac{1}{6} h_3 \rho'''_3 - \frac{1}{4} h'_3 \rho''_3 + \frac{1}{4} h''_3 \rho'_3 - \frac{1}{6} h'''_3 \rho_3 + \frac{4}{3c} w_2 h_3 \rho'_3 - \frac{4}{3c} w_2 h'_3 \rho_3 \\ \delta(\rho_3) h_3 &= -h_2 \rho'_3 + 2h'_2 \rho_3 + \dot{\rho}_3. \end{aligned} \quad (40)$$

Finally, the dynamical equations $E_{1,2} = 0$, obtained from the zero-curvature condition are the following:

$$\partial_+ w_2 = ((c + \alpha_2) \partial^3 + w_2 \partial + \partial w_2) h_2 + (\beta_2 \partial^4 + (w_3 \partial + 2 \partial w_3) h_3, \quad (41a)$$

$$\begin{aligned} \partial_+ w_3 &= (\alpha_3 \partial^3 + 2w_3 \partial + \partial w_3) h_2 \\ &- \frac{1}{12} ((c + \beta_3) \partial^5 + (2 \partial^3 w_2 + 3 \partial^2 w_2 \partial + 3 \partial w_2 \partial^2 + 2 w_2 \partial^3) + \frac{16}{c} w_2 \partial w_2) h_3. \end{aligned} \quad (41b)$$

The α_i 's and β_i 's are linear combinations³ of the κ_i 's. It is remarkable that they only occur in the shift of the central charges, as in the $sl(2, R)$ case. Let us recall that this system leads to the Boussinesq equation by setting $h_2 = 0$ and $h_3 = \frac{1}{12}$. The particular case $\kappa_i = \lambda_i = 0$ reproduces the analysis of Das et al^[7].

The previously mentioned covariant operators $\mathcal{L}_{(2)}$ and $\mathcal{L}_{(3)}$ can be easily seen to correspond to the matrix determinant of the gauge covariant operators $\mathcal{D}_- = \partial + A_-$ respectively for the gauge choice of Eq.(21) and Eq.(33). This observation will be very

³ Notice that the explicit expressions for the α_i 's and the β_i 's are irrelevant in this classical analysis.

useful in the next section where we investigate the general $sl(N, R)$ case. We will adopt the approach of partial gauge fixing *à la* Polyakov in obtaining the w_N algebras in the A_- sector and the corresponding metric fields in the A_+ sector. The guiding principle in the A_- sector is to keep the correspondence between the scalar Lax operator $\mathcal{L}_{(N)}$ and the matrix covariant operator $\mathcal{D}_-$. Although this approach seems lacking in physical motivations, recent works have illustrated connection of this method with 2+1 dimensional Chern-Simons theories^[16]. Hence, assuming validity of the Chern-Simons approach for the general $sl(N, R)$ case, we will proceed with the present analysis.

IV. General Lax pair: from $sl(3, R)$ to $sl(N, R)$

This section presents a general algorithm to calculate the matrix Lax pair associated with $sl(N, R)$. This is done in three steps. First, the A_- sector of the gauge field is fixed in such a way that the remaining degrees of freedom in A_- , the so-called w_i 's, are Virasoro quasi-primary fields under the remaining gauge preserving transformations. The requirement of quasi-primary nature of the w_i 's avoids the complicated non-linearities that would have occurred in fixing the interaction terms in A_+ with strictly primary w_i 's. In the second step, we define the A_+ sector by introducing a new family of fields, the h_i 's, which are analogs of the metric field for the w_N generators. Our objective now is to obtain, as in the $sl(2, R)$ case, the relevant Ward identities and the consistent constraints involving the h_i 's from the zero-curvature condition. This procedure determines most of the components of A_+ ; the remaining freedom is related with the constraint equations themselves (analogs of the κ and λ of the $sl(2, R)$ case) and the symmetry properties of the h_i 's under diffeomorphisms. This leads to the last step, similar to the first one, where we require the fields $h_i, i > 2$ to transform like currents under the fundamental Virasoro symmetry:

$$\delta(\rho)h_{i,i>2} = -(i-1)h_i\rho' + h_i'\rho. \quad (42)$$

For h_2 , we want to maintain Eq.(3), at least for the finite subgroup of diffeomorphism. This requirement and Eq.(42) imply that $\partial^i h_i$ are dimensionless quantities and, hence, the constraint equations

$$\Phi_i \equiv \kappa_i \partial^i h_i - \lambda_i \sim 0, \quad \forall i = 2, 3, \dots, N \quad (43)$$

are meaningful. This gives us the opportunity to introduce N “cosmological constants” λ_i and N parameters κ_i . Vanishing of the κ_i 's and the λ_i 's eliminates these constraints.

We now propose an algorithm that realizes for any $sl(N, R)$ algebra the gauge choice that satisfies the previous conditions. Keeping the form of the gauge suggested by the

$sl(2, R)$ and $sl(3, R)$ cases, let us write

$$A_- = \begin{pmatrix} 0 & -W_2 & -W_3 & -W_4 & \dots & -W_N \\ c_1 & 0 & \dots & \dots & & 0 \\ 0 & 1 & 0 & \dots & & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 \\ \vdots & & & & & 0 \\ 0 & 0 & & & 1 & 0 \end{pmatrix}, \quad (44)$$

$$A_+ = \begin{pmatrix} \dots & \dots & \dots & \dots & \dots \\ H_2 & \dots & \dots & \dots \\ H_3 & \dots & \dots \\ H_4 & \dots \\ \vdots \\ H_N & \dots \end{pmatrix}. \quad (45)$$

We will put $c_1 = 1$ for convenience; this parameter scales the central charge. The other constant components of A_- could have been chosen different from unity; however, rescaling of the primary fields in A_- eliminates these parameters in both the resulting Poisson brackets and the dynamical equations.

The $N - 1$ fields W_i and H_i have been considered in the Refs.[6,7]. It is well-known that the W_i 's are *not* quasi-primary fields and the H_i 's do not transform as definite spin currents. Let us consider the following change of variables:

$$\begin{cases} W_i = \sum_{j=2}^i (-1)^{i-j} A_{ij}(N) \partial^{i-j} w_j, & \text{with } A_{ij}(N) = \frac{\binom{i-1}{i-j} \binom{N-j}{i-j}}{\binom{i+j-1}{i-j}} \\ H_i = \sum_{j=i}^N B_{ji}(N) \partial^{i-j} h_j, & \text{with } B_{ij}(N) = (-1)^{i-j} \frac{\binom{i-1}{i-j} \binom{N-j}{i-j}}{\binom{2i-2}{i-j}}, \end{cases} \quad (46)$$

that we will justify later. The gauge field A_- being thus completely defined, the zero-curvature condition yields the following recurrent relations on the matrix elements a_{ij} of A_+

$$\begin{cases} a_{i,j+1} - a_{i-1,j} = a'_{i,j} - H_i W_j, \\ a_{k,k} = \sum_{i=2}^k (a'_{i,i-1} - H_i W_{i-1}) - \sum_{i=2}^k \frac{N+1-i}{N} (a'_{i,i-1} - H_i W_{i-1}), \\ a_{N-i,N} = -(a'_{N-i+1,N} - H_{N-i+1} W_N). \end{cases} \quad (47)$$

These equations determine all the elements of A_+ in terms of the $N - 1$ fields H_i 's. Let us ignore the constraints for the moment and concentrate on the symmetries. It is rather obvious that the issue of the gauge preserving symmetries $\delta A_- \equiv \partial_- \epsilon + [A_-, \epsilon]$ is similar to

the zero-curvature condition since, on the constant elements of the gauge field in Eq.(44), the gauge preserving condition amounts to solving $\partial_- \epsilon - \partial_+ A_- + [A_-, \epsilon] = 0$. Therefore, the matrix ϵ is determined by its first column

$$\epsilon = \begin{pmatrix} \dots & \dots & \dots & \dots & \dots \\ \epsilon_2 & \dots & \dots & \dots & \\ \epsilon_3 & \dots & \dots & & \\ \epsilon_4 & \dots & & & \\ \vdots & & & & \\ \epsilon_N & \dots & & & \end{pmatrix}, \quad (48)$$

and the recurrent equations (47) can be used to compute all the other components. The knowledge of the ϵ leads to the proper variations of the w_i 's and of the h_i 's. It is remarkable that the generalization of Eq.(27), in order to fulfill the symmetry requirements previously described, is the following redefinition of the gauge parameter:

$$\epsilon_i = \sum_{j=i}^N B_{ji}(N) \partial^{i-j} \rho_j, \quad (49)$$

with the matrix $B(N)$ defined in Eq.(46). More precisely, looking only at the variations induced by ρ_2 , the gauge transformations lead to the following expressions

$$\begin{cases} \delta w_2 = 2 w_2 \rho'_2 + w'_2 \rho_2 + C(N) \rho_2''', \\ \delta w_{2k+1} = (2k+1) w_{2k+1} \rho'_2 + w'_{2k+1} \rho_2 + \sum_{l=1}^{k-1} C_{2k+1}^l(N) \Omega_l^{2k+1}, \\ \delta w_{2k} = 2k w_{2k} \rho'_2 + w'_{2k} \rho_2 + \sum_{l=1}^{k-1} C_{2k}^l(N) \Omega_l^{2k}. \end{cases} \quad (50)$$

We have defined the central charge $C(N) = \frac{(N-1)N(N+1)}{12}$ and

$$\Omega_l^q = \sum_{i=0}^{2(l-1)} \frac{(-1)^i \sigma_i(l)}{\binom{2q-4l+i-1}{i}} w_{q-2l}^{(i)} \rho_2^{(2l-i+1)}, \quad (51)$$

where $\sigma_i(l)$ are given by:

$$\sigma_0(l) = 1, \quad \sigma_1(l) = 2(l-1)(2l+1), \quad \sigma_2(l) = l(l-1)(2l+1)(2l-3),$$

and,

$$\sigma_{i \geq 3}(l) = \frac{\binom{2l+1}{i} \binom{2(l-1)}{i-3} \binom{2l+1-i}{3}}{\binom{i}{3}}. \quad (52)$$

Let us note that the derivatives appearing on the parameter ρ_2 are always greater or equal to three which implies that we have defined the gauge field A_- in terms of quasi-primary fields. Furthermore, the dependance on N , i.e. on the central charge, is completely contained in the coefficients $C_k^l(N)$. A lengthy calculation leads to the following conjecture^[28]

$$C_k^l(N) = \frac{(l-1)!}{2^l(2l+1)} \binom{N-k+2l}{2l} \binom{k-l-2}{l-1} \frac{(2(k-l)-1)lN + (k-l)(k-l-1) + l^2}{\prod_{i=1}^l (2(k-i)-1)} \quad (53)$$

Finally, let us mention that we obtain the correct symmetry variations of the h_i 's for $i > 2$:

$$\delta(\rho_2)h_{i>2} = -(i-1)h_i \rho_2' + h_i' \rho_2.$$

However, the variation of h_2 involves higher derivative terms which vanish only in the finite subgroup of the diffeomorphisms. This point will be illustrated below with some particular cases $N \geq 4$.

The ‘‘cosmological constants’’ can now be added into the A_+ sector. We consider the following shift of the previous gauge field A_+ :

$$A_+ \rightarrow A_+ + \begin{pmatrix} 0 & \lambda_2 - \kappa_2 \partial^2 h_2 & 0 & 0 & \dots & 0 \\ 0 & 0 & \lambda_3 - \kappa_3 \partial^3 h_3 & 0 & \dots & 0 \\ 0 & 0 & 0 & \lambda_4 - \kappa_4 \partial^4 h_4 & \dots & \\ \vdots & & & & & \vdots \\ 0 & \dots & & 0 & \lambda_N - \kappa_N \partial^N h_N & \\ 0 & \dots & & & & 0 \end{pmatrix}. \quad (54)$$

The resulting zero curvature equations exhibit the expected dynamical equations on the fields w_k with, as already noticed in the $N = 2$ and $N = 3$ cases, a shift in the coefficients of the higher derivative of the h_i 's. The diagonal entries of F_{+-} now exhibit the expected $N - 1$ equations $\Phi_i \sim 0$.

It is easy to show that the presence of constraints restricts the gauge preserving symmetry to that of the finite subgroup of the diffeomorphisms. Of course, such restriction is also seen in the quantum regime where the relevance of the constraints is well understood in terms of the ghost system. Let us only mention that, similar to Eq.(5), the operators h_i 's are expanded in terms of $sl(N, R)$ current operators

$$h_i = \sum_{n=2}^{2i-2} x^{-n} J_i^{(n)}(x^+). \quad (55)$$

Such expansions are legitimate by the equations $\partial^{2i-1} h_i = 0$ that generalize the constant curvature equation and are easily obtainable from the dynamical equations with the w_i 's

equal to zero. Then, the equations $\Phi_i \sim 0$ lead to the following system of constraints

$$J_i^{(i)} \sim \text{const}, \quad \text{and} \quad \left\{ \begin{array}{l} J_i^{(i+1)} \sim 0, \\ J_i^{(i+2)} \sim 0, \\ \vdots \\ J_{i,i>2}^{(j)} \sim 0, \\ \vdots \\ J_i^{(2i-2)} \sim 0. \end{array} \right. , i = 2, 3, \dots, N \quad (56)$$

Dimensional analysis of this set of constraints completely describes the weight spectrum of the necessary ghost systems. The resulting central charge contribution is^[10,28] $c_{ghost} = -n^2(n+1)^2 + n(n+1) + 2$.

Let us illustrate the previous results with the case of $N = 4$. The definitions in Eq.(46) give

$$\begin{aligned} W_2 &= w_2, \\ W_3 &= w_3 - w'_2, \\ W_4 &= w_4 - \frac{1}{2}w'_3 + \frac{3}{10}w''_2; \end{aligned} \quad (57)$$

and

$$\begin{aligned} H_2 &= h_2 - h'_3 + \frac{1}{5}h''_4, \\ W_3 &= h_3 - h'_4, \\ W_4 &= h_4. \end{aligned} \quad (58)$$

The gauge is then uniquely determined through the relations (47) and, finally, the shift (54) introduces the cosmological constants. Inspecting the gauge transformations that maintain the gauge so defined, we obtain the $c = 5$ Virasoro algebra for w_2 together with the following infinitesimal variations:

$$\begin{aligned} \delta(\rho_2) w_3 &= 3w_3 \rho'_2 + \rho_2 w'_3, \\ \delta(\rho_2) w_4 &= 4w_4 \rho'_2 + \rho_2 w'_4 + \frac{9}{10}w_2 \rho'''_2, \\ \delta(\rho_2) h_2 &= -h_2 \rho'_2 + \rho_2 h'_2 - \frac{9}{10}h_4 \rho'''_2 + \dot{\rho}_2, \\ \delta(\rho_2) h_3 &= -2h_3 \rho'_2 + \rho_2 h'_3, \\ \delta(\rho_2) h_4 &= -3h_4 \rho'_2 + \rho_2 h'_4, \end{aligned} \quad (59)$$

and the higher spin variations listed in the Appendix A. The variation $\delta(\rho_2) w_4$ is a consequence of our gauge choice linear in the w_i 's. However, we can redefine the parameterisation

of the gauge and introduce a non-linear term in W_4 :

$$W_4 = w_4 - \frac{1}{2}w'_3 + \frac{3}{10}w''_2 + \frac{9}{100}w_2^2 \quad (60)$$

such that w_4 is now primary, like w_3 . This remark leads us to the justification of the parameterisation of W_i in Eq.(46). It mainly relies on a theorem due to Itzykson et al.^[21] in the context of the scalar Lax operator $\mathcal{L}_{(N)}$; the matrix determinant being the operation that connects our approach to theirs. Similar to Ref[21], our gauge is thus parameterized with quasi-primary fields. The previous redefinition (60) illustrates the missing part of the theorem, i.e. the non-linear contribution in W_i that makes *all* the $w_{i>2}$ primary. Using the variations in Eq.(50), the expression of W_4 given by Eq.(60) can be generalised to higher $N \geq 4$:

$$W_4 = w_4 + \sum_{j=2}^3 (-1)^{4-j} A_{4j}(N) \partial^{4-j} w_j + \frac{C_4^1(N)}{C(N)} w_2^2, \quad (61)$$

Further generalizations to higher dimensional W_i 's are certainly possible. For example,

$$\begin{aligned} W_5 &= w_5 + \sum_{j=2}^4 (-1)^{5-j} A_{5j}(N) \partial^{5-j} w_j + \frac{C_5^1(N)}{C(N)} w_2 w_3, \\ W_6 &= w_6 + \sum_{j=2}^5 (-1)^{6-j} A_{6j}(N) \partial^{6-j} w_j + \frac{C_6^1(N)}{C(N)} w_2 w_4 + \frac{C_6^2(N)}{4C(N)} (4w_2 w_2'' - 5w_2'^2) \\ &\quad + \frac{C_6^1(N) C_4^1(N) - 9C_6^2(N)}{6C(N)^2} w_2^3, \\ W_7 &= w_7 + \sum_{j=2}^6 (-1)^{7-j} A_{7j}(N) \partial^{7-j} w_j + \frac{C_7^1(N)}{C(N)} w_2 w_5 + \frac{C_7^2(N)}{21C(N)} (21w_3 w_2'' - 35w_3' w_2') \\ &\quad + 10w_3'' w_2) + \frac{7C_7^1(N) C_5^1(N) - 24C_7^2(N)}{14C(N)^2} w_2^2 w_3, \\ &\text{etc...} \end{aligned} \quad (62)$$

A complete formula for this parameterisation of the W_k 's in terms of w_2 and *primary* w_i 's is under investigation.

The parameterisation of H_i in Eq.(46) is more intriguing since, as shown in Ref.[21], the matrices $A_{ij}(N)$ and $B_{ij}(N)$ satisfy $A.B = 1$. Whereas the fields h_i 's satisfy the symmetry conditions described previously, the Virasoro variation of the metric field h_2 exhibits a contribution that only vanishes on the finite subgroup of the diffeomorphisms. This observation is also true for higher w_N gravities (see Appendix C). However, it is possible to introduce coupling terms in the field H_2

$$H_2 = h_2 - h_3' + \frac{1}{5}(\partial^2 + \frac{9}{2}w_2)h_4 \quad (63)$$

such that the field h_2 now exactly transforms as in Eq.(3).

Let us come back to our $N = 4$ example and consider the zero curvature equations. They exhibit the constraints on the h_i 's and the following dynamical equations

$$\begin{aligned} \partial_+ w_2 = & \left((5 + \alpha_2) \partial^3 + w_2 \partial + \partial w_2 \right) h_2 + \left(\beta_2 \partial^4 + (w_3 \partial + 2 \partial w_3) \right) h_3 \\ & + \left(\gamma_2 \partial^5 + \frac{9}{10} \partial^3 w_2 + (w_4 \partial + 3 \partial w_4) \right) h_4 \end{aligned} \quad (64a)$$

$$\begin{aligned} \partial_+ w_3 = & (\alpha_3 \partial^4 + (2w_3 \partial + \partial w_3)) h_2 - \frac{1}{5} \left((5 + \beta_3) \partial^5 + (2 \partial^3 w_2 + 3 \partial^2 w_2 \partial + 3 \partial w_2 \partial^2 \right. \\ & + 2w_2 \partial^3) - 10(w_4 \partial + \partial w_4) + 5w_2 \partial w_2 \left. \right) h_3 - \frac{1}{10} \left(\gamma_3 \partial^6 + 5 \partial^3 w_3 + 5 \partial^2 w_3 \partial \right. \\ & + 3w_3 \partial^3 \left. \right) + 5 w_2 \partial w_3 \left. \right) h_4 \end{aligned} \quad (64b)$$

and

$$\begin{aligned} \partial_+ w_4 = & \left(\alpha_4 \partial^5 + \frac{9}{10} w_2 \partial^3 + 3w_4 \partial + \partial w_4 \right) h_2 - \frac{1}{10} \left(\beta_4 \partial^6 + \partial^3 w_3 + 3 \partial^2 w_3 \partial + 5 \partial w_3 \partial^2 \right. \\ & + 5w_3 \partial^3 + 5w_3 \partial w_2 \left. \right) h_3 + \frac{1}{100} \left((5 + \gamma_4) \partial^7 + 3 \partial^5 w_2 + 5 \partial^4 w_2 \partial + 6 \partial^3 w_2 \partial^2 \right. \\ & + 6 \partial^2 w_2 \partial^3 + 5 \partial w_2 \partial^4 + 3w_2 \partial^5 + 10(\partial^3 w_4 + 2 \partial^2 w_4 \partial + 2 \partial w_4 \partial^2 + w_4 \partial^3) \\ & + 5(\partial^3 w_2^2 + w_2^2 \partial^3 + 4 \partial w_2 \partial w_2 \partial + 4w_2 \partial^3 w_2) - 75w_3 \partial w_3 + 100(w_2 w_4 \partial + \partial w_2 w_4) \left. \right) h_4 \end{aligned} \quad (64c)$$

We notice that equations (64a,b) with $h_2 = h_3 = 0$, $h_4 = \frac{1}{2}$ and $w_4 = w_2^2$ lead to a coupled KdV system:

$$\begin{cases} \partial_+ w_2 - \frac{1}{2} w_2''' - 3w_2' w_2 = 0, \\ \partial_+ w_3 - \frac{1}{4} w_3''' - \frac{1}{4} w_3' w_2 = 0, \end{cases} \quad (65)$$

whereas Eq.(64c) becomes the following constraint equation:

$$\frac{1}{100} w_2'''' + \frac{1}{20} w_2''' - \frac{3}{5} w_2 w_2''' + \frac{1}{3} w_2^{3'} = \frac{1}{8} w_3^{2'}. \quad (66)$$

V. Conclusion

In this note, we studied the Lax pair formulation of the w_N gravity theories; the analysis being strictly at the classical level. One of the matrix Lax operator $\partial + A_-$ yielded the basic differential operator associated with the n th reduction of the KP hierarchy, after taking the matrix determinant or, equivalently, solving the differential equation $(\partial + A_-)\Psi = 0$ where Ψ takes its value in a jet bundle^[6]. Hence, interestingly, the partial gauge fixing of the A_- sector, designed for obtaining fields transforming as higher spin objects under the residual gauge transformations, turned out to be connected with the process of covariantization of this n th order differential operator. It will be of interest to find out the geometry behind the other members of the $w_N^{(l)}$ algebras. Although these fields were chosen to be quasi-primary for the sake of convenience, the generalisation to primary fields can be achieved by using the above mentioned connection with the scalar covariant operators. We gave a conjectured formula, valid to all orders, for the infinitesimal variations of the w_i 's under diffeomorphism. By studying these variations, it is possible, in principle, to construct the primary fields out of the quasi primary ones. It is of deep interest to obtain an all order formulation for the gauge defined by Eq.(44) in terms of primary fields; this problem is currently under investigation. Furthermore, the work of Zamolodchikov has shown the connection of these covariant operators (in the absence of higher spin fields) with the induced Liouville action, which indicates a large- N analysis of this problem can be carried out using the matrix valued connections thereby clarifying the relationship between the continuum and the lattice model approaches to the two dimensional gravity. The A_+ sector, related with the $sl(N, R)$ current algebra, has been determined modulo scalings of the metric fields h_i 's by constant parameters, by requiring $h_{i>2}$ to transform as currents under the Virasoro symmetry and h_2 to transform as the light-cone metric. The scaling freedom is tied with the level of the Kac-Moody algebra. The relationship of the original H_i 's and the h_i 's is of great interest: it happens to be the inverse of the transformation relating the noncovariant fields in $M_{(N)}$ to the covariant ones in $\mathcal{L}_{(N)}$. The same connection also holds for the gauge parameters; namely the infinitesimal parameters associated with the Virasoro and the higher spin symmetries and the ϵ^- 's appearing in the gauge transformations are connected to each other by an identical relation. We suspect that this intriguing connection originates from the coadjoint orbits of the Lax operator. This point needs further study for clarification.

The other entries in A_+ were determined by demanding that the zero curvature condition should yield the relevant constraints and the Ward identities. It is worth mentioning that the same equations follow from the requirement of the consistency of the gauge fixing. This is natural because the Lie derivative of a gauge field can be described as a gauge transformation if the curvature tensor vanishes. The $F_{+-} = 0$ condition immediately

suggests a connection with the Chern-Simons theories, where similar condition appears as the Gauss law constraint. In fact, this connection has been already illustrated for the w_2 and $w_3^{(2)}$ cases^[16]; the proof for the w_N case is of obvious interest. In light of the beautiful geometry associated with the compact non-abelian Chern-Simons theories, this point needs deeper understanding. Another open question is the construction of the covariant action like that of the KPZ analysis of the $sl(2, R)$ case and understanding of the precise relationship between the non-linear hierarchies appearing in the continuum approach and those of the matrix models.

Acknowledgements

This work was supported in part by funds provided by the Natural Sciences and Engineering Research Council (NSERC) of Canada.

Appendix A:

The infinitesimal higher spin variations of the w_i 's and the h_i 's in the $sl(4, R)$ case are the following:

$$\begin{aligned}
\delta(\rho_3) w_2 &= 3w_3\rho_3' + 2\rho_3 w_3', \\
\delta(\rho_3) w_3 &= 4w_4\rho_3' + 2w_4'\rho_3 + 2w_2\rho_3''' + 3w_2'\rho_3'' + \frac{9}{5}w_2''\rho_3' + \frac{2}{5}w_2'''\rho_3 \\
&\quad - (w_2^2\rho_3' + w_2w_2'\rho_3) - \rho_3''''', \\
\delta(\rho_3) w_4 &= -\frac{7}{5}w_3\rho_3''' - \frac{7}{5}w_3'\rho_3'' - \frac{3}{5}w_3''\rho_3' - \frac{1}{10}w_3'''\rho_3 \\
&\quad - \frac{1}{2}(w_2w_3\rho_3' + w_3'w_2\rho_3), \\
\delta(\rho_3) h_2 &= \frac{2}{5}h_3\rho_3''' - \frac{3}{5}h_3'\rho_3'' + \frac{3}{5}h_3''\rho_3' - \frac{2}{5}h_3'''\rho_3 \\
&\quad - \frac{1}{2}(h_4'w_3\rho_3 + h_4w_3'\rho_3) + h_3w_2\rho_3' - h_3'w_2\rho_3, \\
\delta(\rho_3) h_3 &= 2h_2'\rho_3 - h_2\rho_3' + \dot{\rho}_3 \\
&\quad + \frac{1}{2}h_4\rho_3''' - \frac{1}{2}h_4'\rho_3'' + \frac{3}{10}h_4''\rho_3' - \frac{1}{10}h_4'''\rho_3 \\
&\quad + \frac{1}{2}(h_4'w_2\rho_3 + h_4w_2'\rho_3), \\
\delta(\rho_3) h_4 &= 2h_3'\rho_3 - 2h_3\rho_3',
\end{aligned}$$

$$\begin{aligned}
\delta(\rho_4) w_2 &= 4w_4\rho_4' + 3\rho_4 w_4' + \frac{9}{10}(w_2\rho_4''' + 3w_2'\rho_4'' + 3w_2''\rho_4' + 3w_2'''\rho_4), \\
\delta(\rho_4) w_3 &= -\frac{1}{10}(14w_3\rho_4''' + 28w_3'\rho_4'' + 20w_3''\rho_4' + 5w_3'''\rho_4) - \frac{1}{2}(w_2w_3\rho_4' + w_2w_3'\rho_4), \\
\delta(\rho_4) w_4 &= \frac{1}{10}(w_4\rho_4''' + 9w_4'\rho_4'' + 5w_4''\rho_4' + 6w_4'''\rho_4) \\
&\quad + \frac{7}{25}w_2\rho_4'''' + \frac{7}{10}w_2'\rho_4''' + \frac{21}{25}w_2''\rho_4'' + \frac{14}{25}w_2'''\rho_4' + \frac{1}{5}w_2''''\rho_4 + \frac{3}{100}w_2'''''\rho_4 \\
&\quad - \frac{3}{4}(w_3^2\rho_4' + w_3w_3'\rho_4) + 2w_4w_2\rho_4' + w_4'w_2\rho_4 + w_4w_2'\rho_4 + \frac{1}{2}w_2^2\rho_4''' \\
&\quad + \frac{3}{2}w_2w_2'\rho_4'' + \frac{1}{2}w_2'^2\rho_4' + \frac{11}{10}w_2w_2''\rho_4' \\
&\quad + \frac{3}{10}w_2'w_2''\rho_4 + \frac{3}{10}w_2w_2'''\rho_4 + \frac{1}{20}\rho_4''''', \\
\delta(\rho_4) h_2 &= \frac{9}{10}h_2'''\rho_4 - \frac{3}{100}h_4\rho_4'''' + \frac{1}{20}h_4'\rho_4''' - \frac{3}{50}h_4''\rho_4'' + \frac{3}{50}h_4'''\rho_4' - \frac{1}{20}h_4''''\rho_4' \\
&\quad + \frac{3}{100}h_4'''''\rho_4 - \frac{3}{10}w_2h_4\rho_4''' - \frac{3}{5}w_2'h_4\rho_4'' + \frac{1}{5}w_2h_4'\rho_4' - \frac{1}{5}w_2h_4''\rho_4 - \frac{3}{5}w_2''h_4\rho_4' \\
&\quad + \frac{3}{5}w_2'h_4''\rho_4 + \frac{3}{10}w_2h_4'''\rho_4 + w_4h_4\rho_4' + w_4'h_4\rho_4 + \frac{1}{2}h_3w_3\rho_4 + \frac{1}{2}h_3w_3'\rho_4, \\
\delta(\rho_4) h_3 &= \frac{1}{10}(h_3\rho_4''' - 3h_3'\rho_4'' + 5h_3''\rho_4' - 5h_3'''\rho_4) \\
&\quad - \frac{1}{2}w_2h_3\rho_4 - \frac{1}{2}w_2'h_3\rho_4 - \frac{3}{4}w_3h_4\rho_4 + \frac{3}{4}w_3h_4'\rho_4, \\
\delta(\rho_4) h_4 &= 3h_2'\rho_4 - h_2\rho_4' + \dot{\rho}_4 \\
&\quad - \frac{1}{10}(h_4\rho_4''' - 2h_4'\rho_4'' - 2h_4''\rho_4' + h_4'''\rho_4 + h_4'w_2\rho_4 - h_4w_2\rho_4').
\end{aligned}$$

Appendix B:

Looking at the gauge symmetries preserving the gauge as defined in Eq.(44) for $sl(10, R)$, we get the following Virasoro transformations of the w_i 's, $2 \leq i \leq 10$:

$$\begin{aligned}
\delta(\rho_2)w_2 &= 2w_2\rho_2' + w_2'\rho_2 + \frac{165}{2}\rho_2''', \\
\delta(\rho_2)w_3 &= 3w_3\rho_2' + w_3'\rho_2, \\
\delta(\rho_2)w_4 &= 4w_4\rho_2' + w_4'\rho_2 + \frac{266}{5}w_2\rho_2''',
\end{aligned}$$

$$\begin{aligned}
\delta(\rho_2)w_5 &= 5w_5\rho_2' + w_5'\rho_2 + \frac{83}{2}w_3\rho_2''', \\
\delta(\rho_2)w_6 &= 6w_6\rho_2' + w_6'\rho_2 + \frac{185}{6}w_4\rho_2'''' \\
&\quad + \frac{52}{3}w_2\rho_2''''' - \frac{130}{3}w_2'\rho_2'''' + \frac{130}{3}w_2''\rho_2''', \\
\delta(\rho_2)w_7 &= 7w_7\rho_2' + w_7'\rho_2 + \frac{235}{11}w_5\rho_2''' \\
&\quad + \frac{119}{11}w_3\rho_2'''' - \frac{595}{33}w_3'\rho_2'''' + \frac{170}{11}w_3''\rho_2''', \\
\delta(\rho_2)w_8 &= 8w_8\rho_2' + w_8'\rho_2 + \frac{173}{13}w_6\rho_2''' \\
&\quad + \frac{762}{143}w_4\rho_2'''' - \frac{1905}{286}w_4'\rho_2'''' + \frac{635}{429}w_4''\rho_2''' \\
&\quad + \frac{23}{33}w_2\rho_2''''' - \frac{161}{33}w_2'\rho_2''''' + \frac{483}{55}w_2''\rho_2'''' - \frac{161}{33}w_2'''\rho_2'''' + \frac{23}{33}w_2''''\rho_2''', \\
\delta(\rho_2)w_9 &= 9w_9\rho_2' + w_9'\rho_2 + \frac{173}{13}w_7\rho_2''' \\
&\quad + \frac{762}{143}w_5\rho_2'''' - \frac{1905}{286}w_5'\rho_2'''' + \frac{635}{429}w_5''\rho_2''' \\
&\quad + \frac{23}{33}w_3\rho_2''''' - \frac{161}{33}w_3'\rho_2''''' + \frac{483}{55}w_3''\rho_2'''' - \frac{161}{33}w_3'''\rho_2'''' + \frac{23}{33}w_3''''\rho_2''', \\
\delta(\rho_2)w_{10} &= 10w_{10}\rho_2' + w_{10}'\rho_2 + \frac{173}{13}w_8\rho_2''' \\
&\quad + \frac{762}{143}w_6\rho_2'''' - \frac{1905}{286}w_6'\rho_2'''' + \frac{635}{429}w_6''\rho_2''' \\
&\quad + \frac{23}{33}w_4\rho_2''''' - \frac{161}{33}w_4'\rho_2''''' + \frac{483}{55}w_4''\rho_2'''' - \frac{161}{33}w_4'''\rho_2'''' + \frac{23}{33}w_4''''\rho_2''' \\
&\quad + \frac{23}{33}w_2\rho_2'''''' - \frac{161}{33}w_2'\rho_2'''''' + \frac{483}{55}w_2''\rho_2''''' - \frac{161}{33}w_2'''\rho_2''''' + \frac{23}{33}w_2''''\rho_2'''' \\
&\quad - \frac{161}{33}w_2''''\rho_2'''' + \frac{23}{33}w_2''''' \rho_2'''.
\end{aligned}$$

Appendix C:

As mentioned in the text, the Virasoro variation of h_i with $i > 2$ is that of a current with dimension i ,

$$\delta(\rho_2)h_{i,i>2} = -(i-1)h_i\rho_2' + h_i'\rho_2.$$

However, the variation of h_2 involves higher derivative terms for $N \geq 4$. Denoting

$$\mathcal{L}_{\rho_2} h_2 = -h_2 \rho_2' + \rho_2 h_2' + \dot{\rho}_2,$$

we obtain for $N \geq 4$

$$\delta(\rho_2) h_2 = \mathcal{L}_{\rho_2} h_2 + \Gamma_N,$$

where Γ_N is vanishing in the finite subgroup of diffeomorphism, in which case $\rho_2''' = 0$. In addition to $\delta(\rho_2) h_2$ given in Eq.(59), some explicit expressions are given below

$$\begin{aligned} \Gamma_5 &= -\frac{16}{5} h_4 \rho_2''', \\ \Gamma_6 &= -\frac{37}{5} h_4 \rho_2''' - \frac{5}{7} h_6 \rho_2'''' - \frac{5}{7} h_6' \rho_2'''' - \frac{10}{63} h_6'' \rho_2''', \\ \Gamma_7 &= -14 h_4 \rho_2''' - \frac{57}{14} h_6 \rho_2'''' - \frac{57}{14} h_6' \rho_2'''' - \frac{19}{21} h_6'' \rho_2''', \\ \Gamma_8 &= -\frac{47}{2} h_4 \rho_2''' - \frac{96}{7} h_6 \rho_2'''' - \frac{96}{7} h_6' \rho_2'''' - \frac{64}{21} h_6'' \rho_2''' \\ &\quad - \frac{7}{12} h_8 \rho_2'''' - \frac{7}{6} h_8' \rho_2'''' - \frac{21}{26} h_8'' \rho_2'''' - \frac{35}{156} h_8''' \rho_2'''' - \frac{35}{1716} h_8'''' \rho_2''', \\ \Gamma_9 &= -\frac{182}{5} h_4 \rho_2''' - \frac{71}{2} h_6 \rho_2'''' - \frac{71}{2} h_6' \rho_2'''' - \frac{71}{9} h_6'' \rho_2''' \\ &\quad - \frac{68}{15} h_8 \rho_2'''' - \frac{136}{15} h_8' \rho_2'''' - \frac{408}{65} h_8'' \rho_2'''' - \frac{68}{39} h_8''' \rho_2'''' - \frac{68}{429} h_8'''' \rho_2''', \\ \Gamma_{10} &= -\frac{256}{5} h_4 \rho_2''' - 78 h_6 \rho_2'''' - 78 h_6' \rho_2'''' - \frac{52}{3} h_6'' \rho_2''' \\ &\quad - \frac{299}{15} h_8 \rho_2'''' - \frac{598}{15} h_8' \rho_2'''' - \frac{138}{5} h_8'' \rho_2'''' - \frac{23}{3} h_8''' \rho_2'''' - \frac{23}{33} h_8'''' \rho_2''' \\ &\quad - \frac{27}{55} h_{10} \rho_2'''' - \frac{81}{35} h_{10}' \rho_2'''' - \frac{324}{187} h_{10}'' \rho_2'''' - \frac{189}{187} h_{10}''' \rho_2'''' \\ &\quad - \frac{567}{1870} h_{10}'''' \rho_2'''' - \frac{81}{1870} h_{10}'''' \rho_2'''' - \frac{27}{12155} h_{10}'''' \rho_2'''. \end{aligned}$$

We notice that Γ_N involves only even dimensional h_i 's, e.g. $h_4, h_6, \dots$. As already mentioned in the $sl(4, R)$ case, these higher derivative terms can be removed by a redefinition of H_2 that introduces coupling terms. For example,

$$\begin{aligned} N=5 : H_2 &= h_2 - \frac{3}{2} h_3' + \frac{3}{5} (\partial^2 - \frac{8}{15} w_2) h_4 - \frac{1}{14} h_5''', \\ N=6 : H_2 &= h_2 - 2h_3' + \frac{6}{5} (\partial^2 - \frac{37}{105} w_2) h_4 - \frac{2}{7} h_5''' \\ &\quad + \frac{1}{42} (\partial^4 - \frac{12}{7} (\frac{2}{9} w_2 \partial^2 + w_2' \partial + w_2'' - 4w_2^2)) h_6, \\ N=7 : H_2 &= h_2 - \frac{5}{2} h_3' + 2(\partial^2 - \frac{1}{4} w_2) h_4 - \frac{5}{7} h_5''' \\ &\quad + \frac{5}{42} (\partial^4 + \frac{171}{140} (\frac{2}{9} w_2 \partial^2 + w_2' \partial + w_2'' - 4w_2^2)) h_6 - \frac{1}{132} h_7''''', \end{aligned}$$

will lead to $\delta(\rho_2) h_2 = \mathcal{L}_{\rho_2} h_2$. A general all order expression for H_2 is under investigation.

References

- [1] A.M. Polyakov, Mod. Phys. Lett. **A2** (1987) 893; See also *Two-Dimensional quantum gravity and Superconductivity at high T_c* in Fields, Strings and Critical Phenomena, eds. E. Brézin and J. Zinn-Justin, Les Houches 1988, Elsevier Science Publishers, Amsterdam, (1989).
- [2] A.H. Chamseddine and M. Reuter, Nucl. Phys. **B317** (1989) 757;
 A. Alekseev and S. Shatashvili, Nucl. Phys. **B323** (1989) 719;
 M. Bershadsky and H. Ooguri, Comm. Math. Phys. **126** (1989) 49;
 A.M. Polyakov, Mod. Phys. Lett. **A6** (1991) 635;
 K. Isler and J-M. Lina, Nucl. Phys. **B358** (1991) 713;
 D. Kutasov, E. Martinec and N. Seiberg, preprint PUPT- 1923, RU-91-49 and references therein.
 For the approach in the conformal gauge, see E. D'Hoker, *Lecture notes on 2-d quantum gravity and Liouville theory*, preprint UCLA/91/TEP/35 (1991), and references therein.
- [3] M.R. Douglas, Phys. Lett. B238 (1990) 176;
 M.A. Awada and Z. Qiu, Phys. Lett. B245 (1990) 85;
 M.A. Awada and S.J. Sin, Institute of fundamental theory preprint UFIFT HEP-91-3 (1991).
- [4] V.G. Knizhnik, A.M. Polyakov and A.B. Zamolodchikov, Mod. Phys. Lett. **A3** (1988) 819;
 J. Distler and H. Kawai, Nucl. Phys. **B321** (1989) 509.
- [5] A.M. Polyakov, Int. J. Mod. Phys. **A5** (1990) 833.
- [6] A. Bilal, V.V. Fock and I.I. Kogan, Nucl. Phys. **B359** (1991) 635, and references therein;
 H. Ooguri, K. Schoutens, A. Sevrin and P. Van Nieuwenhuizen, Stony Brook preprint ITP-SB-91-16, RIM-764.
 Other approaches to w_N gravity can be found in the following papers
 C.M. Hull, Phys. Lett. B240 (1990) 110; *ibid.* B269 (1991) 257;
 K. Schoutens, A. Sevrin and P. Van Nieuwenhuizen, Stony Brook preprint ITP-SB-91-21; Phys. Lett. B243 (1990) 295.
 R. Parthasarathy and K.S. Viswanathan, Simon Fraser Univ. preprint (1991).
- [7] A. Das, W-J. Huang and S. Roy, Rochester preprint UR1197, ER13065-652.
- [8] A.M. Polyakov and P. Wiegmann, Phys. Lett. B131 (1983) 121; Phys. Lett. B141 (1984) 223.

- [9] J.M. Lina and P.K. Panigrahi, Mod. Phys. Lett. **A6** (1991) 3517.
- [10] A. Gerasimov, A. Levin and A. Marshakov, Nucl. Phys. **B317** (1989) 757.
- [11] A.B. Zamolodchikov, Teor. Mat. Fiz 65 (1985) 1205;
V.A. Fateev and S.L. Lukyanov, Int. Jour. Mod. Phys. **A3** (1988) 507.
- [12] A. Bilal, *Introduction to W-algebras* CERN-TH. preprint 6083/91, LPTENS 91/10 (1991).
- [13] J. Balog, L. Fehér, L. O’Raifeartaigh and A. Wipf, Phys. Lett. B227 (1989) 214;
L. Fehér, L. O’Raifeartaigh, P. Ruelle, I. Tsusui and A. Wipf, preprint U.de M.-LPN-TH.71, DIAS-STP-91-29 (1991) and references therein.
- [14] F. Bais, P. Bouwknegt, M. Surridge and K. Schoutens, Nucl. Phys. **B304** (1988) 348, *ibid.* 371.
- [15] A. Bilal and J.L. Gervais, Nucl. Phys. **B314** (1989) 646, *ibid.* **B318** (1989) 579;
- [16] H. Verlinde, Nucl. Phys. **B337** (1990) 652;
A. Bilal, Phys. Lett. B267 (1991) 487.
- [17] K. Yamagishi, Phys. Lett. B205 (1988) 466;
P. Mathieu, Phys. Lett. B208 (1988) 101;
Q. Wang, P.K. Panigrahi, U. Sukhatme and W-Y. Keung, Nucl. Phys. **B344** (1990) 196;
I. Bakas, Phys. Lett. B228 (1989) 57.
- [18] J.L. Gervais, Phys. Lett. B160 (1985) 277;
F. Magri, Jour. of Math. Phys. 19 (1978) 1156.
See also P.J. Olver, *Applications of Lie groups to differential equations*, Springer, Berlin (1986);
A. Das, *Integrable Models*, World Scientific (1989).
- [19] W. Scherer, Lett. Math. Phys. **17** (1989) 45.
- [20] A. Das and S. Roy, Rochester preprint UR-1151 (1990).
- [21] B.L. Feigin and D.B. Fuchs, Func. Anal. & Appl. **17** (1982) 114;
P. Di Francesco, C. Itzykson and J-B. Zuber, Princeton preprint SPhT/90-149, PUPT-1211.
- [22] A.B. Zamolodchikov, preprint ITEP 89-84 (1989).
- [23] Y. Matsuo, Nucl. Phys. **B342** (1990) 643.
- [24] V. Drinfeld and V. Sokolov, Jour. Sov. Math. 30 (1985) 1975;
I.M. Gelfand and L.A. Dikki, Funct. Anal. Appl. **10** (1976) 259;

- A.M. Adler, Invent. Math. **50** (1979) 219.
- [25] A.M. Adler, see Ref.[24].
- [26] K. Isler and C.A. Trugenberger, Phys. Rev. Lett. 63 (1989) 834;
E. D'Hoker, preprint UCLA/91/TEP/8.
- [27] This analysis has been made possible by the use of symbolic programing with *Mathematica*. Formula (53) has been checked up to $N = 40$, using a program available by e-mail from the following adress: *lina@lps.umontreal.ca*.
- [28] J. Pawelczyk, preprint WIS-90/28/june (1990).